JOURNAL OF PALEONTOLOGY

INDEX TO VOLUME 23

AUTHOR INDEX

		No.	Page
1.	ASANO, KIYOSHI, Foraminifera from the Asagai formation (Tertiary) of Fukushima prefecture, Japan	5	473
2.	ASANO, KIYOSHI, The foraminiferal genus *Cruciloculina* d'Orbigny, 1839	5	479
3.	ASANO, KIYOSHI, New Miocene Foraminifera from Japan	4	423
4.	BANDY, ORVILLE L., New names for two species of Foraminifera from Little Stave Creek, Clarke County, Alabama	4	440
5.	BARTENSTEIN, H., and BRAND, E., New genera of Foraminifera from the Lower Cretaceous of Germany and England	6	669
6.	BOCK, WILHELM, Triassic chimaeroid egg capsules from the Connecticut Valley	5	515
7.	BRADY, L. F., *Oniscoidichnus*, new name for *Isopodichnus* Brady 1947 not Bornemann 1889	5	573
	BRAND, E., see BARTENSTEIN, H. (5)		
8.	BRANSON, CARL C., Brachiopoda and Mollusca from the *Productus* limestones of the Salt Range, by F. R. Cowper Reed (Review of)	2	230
	BUMP, JAMES D., see WELLES, S. P. (80)		
9.	BURMA, BENJAMIN H., Studies in quantitative paleontology. II. Multivariate analysis—a new analytical tool for paleontology and geology	1	95
10.	DE CIZANCOURT, MARIA, and FRIZZELL, DON L., *Ferayina* in the middle Eocene of Venezuela (Foraminifera, Rotaliidae, Chapmanininae)	5	496
11.	COLE, W. STORRS, Upper Eocene large Foraminifera from the Panama Canal Zone	3	267
12.	COOKE, C. WYTHE, Two Cretaceous echinoids from Peru	1	84
13.	CROUCH, ROBERT W., Pliocene Ostracoda from southern California	6	594
14.	TEN DAM, A., Foraminifera in the Gullmar Fjord and the Skagerak, by Hans Höglund (Review of)	2	231
15.	TEN DAM, A., Repartition stratigraphique du genre *Arenobulimina* Cushman, by N. Grekoff (Review of)	2	231
16.	TEN DAM, A., The Swedish Paleocene and its foraminiferal fauna, by Fritz Brotzen (Review of)	2	232
17.	DECKER, CHARLES E., Five new varieties of graptolites from Tennessee and Virginia	2	214
	DOWNS, H. R., see MILLER, A. K. (47).		
	DOWNS, R. H., see YOUNGQUIST, WALTER (91)		
	DUNCAN, DONALD, see LOCHMAN, CHRISTINA (40)		
18.	DURHAM, J. WYATT, Age of the Carbonera formation near Cúcuta, Colombia	2	146
19.	DUSENBURY, ARTHUR N., The *Hannatoma* fauna in Colombia and Venezuela	2	147
20.	EVITT, WILLIAM R., Stereophotography as a tool of the paleontologist	5	566
	FERRIS, BERNARD J., see JOHNSON, J. HARLAN (38)		
21.	FLOWER, ROUSSEAU H., New genera of Devonian nautiloids	1	74
22.	FREDERICKSON, E. A., Trilobite fauna of the upper Cambrian Honey Creek formation	4	341
23.	FRIZZELL, DON L., Rotalid Foraminifera of the Chapmanininae: their natural distinction and parallelism to the *Dictyoconus* lineage	5	481
24.	FRIZZELL, DON L., and KEEN, MYRA A., On the nomenclature and generic position of *Nautilus beccari* Linné (Foraminifera, "Rotaliidae")	1	106
	FRIZZELL, DON L., see DE CIZANCOURT, MARIA (10)		
25.	GILL, EDMUND D., *Prosopon*, a term proposed to replace the biologically erroneous term *ornament*	5	572
26.	GRIFFITTS, MARY O., A new rudistid from the Niobrara of Colorado	5	471
27.	HAAS, OTTO, Possibility of synonymous homonyms	5	573
28.	HAAS, OTTO, Species without a genus?	1	106
29.	HANNA, G. DALLAS, and HERTLEIN, LEO GEORGE, Two new species of gastropods from the middle Eocene of California	4	392
30.	HANZAWA, SHOSHIRO, A new type of the fusulinid Foraminifera from central Japan	2	205
31.	HATAI, KOTORA, and NISIYAMA, SYÔZÔ, New Tertiary Mollusca from Japan	1	87
32.	HEDBERG, HOLLIS D., The *Hannatoma* fauna in Colombia and Venezuela	2	149
33.	HEMMING, FRANCIS, Revision of the international rules of zoological nomenclature	2	225
	HERTLEIN, LEO GEORGE, see HANNA, G. DALLAS (29)		

34. HOFKER, J., On *Hagenewella* and a new species 4 431
35. HOUGH, JEAN, The subspecies of *Hoplophoneus:* a statistical study 5 536
36. HUSSEY, KEITH M., Louisiana Cane River Eocene Foraminifera 2 109
37. JELETZKY, J. A., *Scaphites morrowi*, new name for *Scaphites pygmaeus* Morrow 1935, non Holzapfel 1888 ..... 3 330
38. JOHNSON, J. HARLAN, and FERRIS, BERNARD J., Tertiary coralline Algae from the Dutch East Indies 2 193
KEEN, A. MYRA, see FRIZZELL, DON L. (24)
39. KEHRER, L., Note on the stratigraphic position of the *Hannatoma* zone in the western Táchira region, Venezuela 2 151
40. LOCHMAN, CHRISTINA, and DUNCAN, DONALD, Homonyms and synonyms in "Early upper Cambrian faunas of central Montana 4 439
41. LOEBLICH, ALFRED R., JR., and TAPPAN, HELEN, Foraminifera from the Walnut formation (Lower Cretaceous) of northern Texas and southern Oklahoma 3 245
42. MCLAREN, D. J., and SUTHERLAND, P. K., *Lithostrotion* from northeast British Columbia and its bearing on the genomorph concept 6 625
43. MANDRA, YORK T., A new species of *Mytilus* from the Pliocene of Humboldt County, California 1 104
44. MARKS, JAY GLENN, Age of the *Hannatoma* fauna 2 153
45. MARKS, JAY GLENN, Nomenclatural units and tropical American Miocene species of the gastropod family Cancellariidae 5 453
46. MILLER, A. K., A giant scaphopod from the Pennsylvanian of Texas 4 387
47. MILLER, A. K., DOWNS, H. R., and YOUNGQUIST, WALTER, Some Mississippian cephalopods from central and western United States 6 600
48. MILLER, A. K., and YOUNGQUIST, WALTER, The Maquoketa coquina of cephalopods .. 2 199
MILLER, A. K., see THOMPSON, M. L. (72), YOUNGQUIST, WALTER (92)
49. MILLER, ROBERT L., An application of the analysis of variance to paleontology 6 635
50. MORISHIMA, MASAO, A new Miocene *Lepidocyclina* from Shizuoka prefecture, Japan .. 2 210
51. NEWELL, NORMAN D., Comments on paleogeography and nomenclature 2 220
NISIYAMA, SYÔZÔ, see HATAI, KOTORA (31)
52. PARKER, PIERRE, Fossil and Recent species of the Pelecypod genera *Chione* and *Securella* from the Pacific Coast 6 577
PATTERSON, S. H., see YOUNGQUIST, WALTER (93)
53. RAO, L. RAMA, Paleontologist—biologist and/or geologist? 6 677
54. RAW, FRANK, Facial sutures in the (hypoparian) trilobites *Loganopeltoides* and *Loganopeltis*, and the validity of these genera 5 510
55. RUSSELL, LORIS S., The relationships of the Alberta Cretaceous dinosaur "*Laosaurus*" *minimus* Gilmore 5 518
56. SINCLAIR, G. WINSTON, Footnotes to *Eobronteus* 5 571
57. SINCLAIR, G. WINSTON, The Ordovician trilobite *Eobronteus* 1 45
58. SINCLAIR, G. WINSTON, *Raymondella, Bancroftina, Reedaspis* 4 438
59. SOHN, I. G., Hinge mechanism of Paleozoic ostracodes 6 691
60. STAINFORTH, R. M., Foraminifera in the upper Tertiary of Egypt 4 419
61. STAINFORTH, R. M., The *Hannatoma* fauna in the Zapotal sands of southwest Ecuador. 2 155
62. STAINFORTH, R. M., Nomenclatural notes on *Pullenia* and *Cibicides* 4 436
63. STONE, BENTON, Age of the Chira group, northwestern Peru 2 156
64. STONE, BENTON, New Foraminifera from northwestern Peru 1 81
65. STUMM, ERWIN C., Three new Devonian species of *Microcyclus* from Michigan and Ontario 5 507
66. STURGEON, MYRON T., and YOUNGQUIST, WALTER, Allegheny conodonts from eastern Ohio 4 380
SUTHERLAND, P. K., see MCLAREN, D. J. (42)
67. SWAIN, FREDERICK M., Early Tertiary Ostracoda from the western interior United States 2 172
68. SWARTZ, FRANK M., Muscle marks, hinge and overlap features, and classification of some Leperditiidae 3 306
TAPPAN, HELEN, see LOEBLICH, ALFRED R., JR. (41)
69. THALMANN, HANS E., Bibliography and index to Foraminifera (Supplements and corrections for the period 1931 to 1947) 5 498
70. THALMANN, HANS E., Bibliography and index to new genera, species, and varieties of Foraminifera for the year 1947 ..... 4 395
71. THALMANN, HANS E., Bibliography and index to new genera, species, and varieties of Foraminifera for the year 1948 ..... 6 641
72. THOMPSON, M. L., The Permian fusulinids of Timor 2 182
73. THOMPSON, M. L., and MILLER, A. K., Permian fusulinids and cephalopods from the vicinity of the Maracaibo Basin in northern South America 1 1

74. TROMP, S. W., The determination of the Cretaceous-Eocene boundary by means of quantitative, generic, microfaunal determinations and the conception "Danian" in the Near East 6 673
75. TROMP, S. W., The value of Globigerinidae ratios in stratigraphy 2 223
76. UNKLESBAY, A. G., An unusual specimen of the genus *Athyris* 3 328
77. WARTHIN, A. S., Lower Ordovician Ostracods of the Siljan district, Sweden, by Ivar Hessland (Review of) 5 576
78. WEINGEIST, LEO, The ostracode genus *Eucytherura* and its species from the Cretaceous and Tertiary of the Gulf Coast 4 364
79. WELLER, J. MARVIN, Paleontologic classification 6 680
80. WELLES, S. P. and BUMP, JAMES D., *Alzadasaurus pembertoni*, a new elasmosaur from the Upper Cretaceous of South Dakota 5 521
81. WILLARD, BRADFORD, An Eden faunule in New Jersey 2 218
82. WILSON, JAMES L., The trilobite fauna of the *Elvinia* zone in the basal Wilberns limestone of Texas 1 25
83. WOOD, ALBERT E., Small mammals from the uppermost Eocene (Duchesnian) near Badwater, Wyoming 5 556
84. YAKOLEV, N. N., *Jaekelicrinus bashkiricus* Yakovlev, n. gen., n. sp. 4 435
85. YEN, TENG-CHIEN, Corrections of fossil localities 3 329
86. YEN, TENG-CHIEN, A new name for *Carinorbis* Yen 5 573
87. YEN, TENG-CHIEN, Review of the Lower Cretaceous fresh-water molluscan faunas of North America 5 465
88. YENNE, KEITH A., *Pedioceras*, a synonym of *Crioceras* (*Pseudocrioceras*) 6 623
89. YOUNGQUIST, WALTER, The cephalopod fauna of the White Pine shale of Nevada 3 276
90. YOUNGQUIST, WALTER, The cephalopod fauna of the White Pine shale of Nevada: supplement 6 613
91. YOUNGQUIST, WALTER, and DOWNS, R. H., Additional conodonts from the Pennsylvanian of Iowa 2 161
92. YOUNGQUIST, WALTER, and MILLER, A. K., Conodonts from the Late Mississippian Pella beds of south-central Iowa 6 617
93. YOUNGQUIST, WALTER, and PATTERSON, S. H., Conodonts from the lower Mississippian Prospect Hill sandstone of Iowa 1 57
YOUNGQUIST, WALTER, see MILLER, A. K. (47, 48), STURGEON, MYRON T. (66)
94. ANONYMOUS, The Center of Paleontological Studies and Documentation of the Paris Museum (C.E.P.D.) 3 340
95. ANONYMOUS, Fellowship administered by the National Research Council 1 108
96. ANONYMOUS, Fellowships and assistantships 3 340
97. ANONYMOUS, The age of the *Hannatoma* mollusk fauna of South America, a symposium: Introduction 2 145

SUBJECT INDEX

Alabama: Foraminifera, new names (4).
Alberta: "*Laosaurus*" *minimus*, Cretaceous, relation hips (55).
Algae, coralline: Tertiary, Dutch East Indies (38).
Allegheny formation: conodonts, Ohio (66).
Alzadasaurus pembertoni, new elasmosaur, Upper Cretaceous, South Dakota (80).
America, tropical: Cancellariidae, Miocene (45).
Ammonites: *Pedioceras* (88); *Scaphites morrowi*, new name (37).
Ammonoidea: *Pedioceras* (88); Permian, Colombia (73); *Scaphites morrowi*, new name (37).
Analysis: multivariate (9); of variance, application to paleontology (49).
Arapahoia convexa, *A. prolixa*, *A. neihartensis*, *A. walcottae* (40).
Arbuckle Mountains: trilobites, upper Cambrian (22).
Arkansas: cephalopods, Mississippian (47).
Arizona: cephalopods, Mississippian (47).
Asagai formation (Tertiary): Foraminifera, Japan (1).
Assistantships: Geology Department, School of Mines and Metallurgy, University of Missouri (96).
Athyris, unusual specimen, Mississippian, Missouri (76).
Bibliography, Foraminifera: new genera, species, and varieties, 1947 (70); 1948 (71); supplements and corrections, 1931–1947 (69); 1939–1947 (71).
Polivinia astoriensis Rau (No. 4, p. 441).
Borneo: coralline algae, Tertiary (38).
Brachiopoda: *Athyris*, Mississippian, Missouri (76); *Bancroftina* Sinclair 1946 (58).
British Columbia: *Lithostrotion*, Mississippian, bearing on genomorph concept (42).
California: *Chione* and *Securella*, Tertiary and Recent (52); Gastropoda, new, Middle Eocene (29); *Hannatoma* fauna, age of (44); *Mytilus*, new species, Pliocene (43); Ostracoda, Pliocene (13).

Cambrian, upper: trilobites, *Elvinia* zone, Texas (82), homonyms and synonyms, Montana (40), Oklahoma (22).
Canada: "*Laosaurus*" minimus, Cretaceous, Alberta, relationships (55); *Microcyclus*, new species, Devonian, Ontario (65); trilobites (hypoparian), Ordovician, Quebec (53).
Canal Zone, Panama: larger Foraminifera, upper Eocene, Gatun Lake (11).
Cancellariidae: nomenclatural units and tropical American Miocene species (45).
Cane River formation (Eocene): Foraminifera, Louisiana, and correlation with Texas and Mississippi formations (36).
Carbonera formation: evidence of upper Eocene age (Colombia) (18).
Carinorbis Yen (86).
Center of Paleontological Studies and Documentation, Paris Museum (94).
Cephalopoda: Mississippian, central and western United States (47); Mississippian, late, eastern Nevada (89, 90); Ordovician, Iowa (48); Permian nautiloids and ammonoids, Colombia (73); *Scaphites morrowi*, new name (37).
Chimaeroid fishes: egg capsules, Triassic, Connecticut Valley (6).
Chione cathcartensis, *C. columbiensis*, *C. fluctifraga*, *C. mathewsonii*, *C. whitneyi* (52).
Chira group, age of, Peru (63).
Cibicides, nomenclatural notes (62).
Classification, paleontologic: suggestions for revision of (79).
Colombia: Cancellariidae, Miocene (45); Carbonera formation, Middle Tertiary age of (18); cephalopods and fusilinids, Permian (73); *Hannatoma* fauna, Tertiary (19, 32, 97); *Pedioceras*, synonym of *Crioceras* (*Pseudocrioceras*) (88).
Colorado: new rudistid, Upper Cretaceous (26); Ostracoda, Eocene and Paleocene (67).
Connecticut: chimaeroid egg capsules, Triassic (6).
Conodonts: Mississippian, late, Iowa (92); Mississippian, lower, southeastern Iowa (93); Pennsylvanian, eastern Ohio (66); Pennsylvanian, early, Iowa (91).
Coosia grandis (40).
Coquina, cephalopod, Ordovician, Iowa (48).
Corals: Devonian, new species, Michigan and Ontario (65); *Lithostrotion*, Mississippian, British Columbia, bearing on genomorph concept (42).
Costa Rica: Cancellariidae, Miocene (45).
Cottaldia australis (12).
Cretaceous: echinoids, Peru (12); -Eocene boundary, established by quantitative generic analysis, Near East (74); *Eucytherura*, Gulf Coast (78); "*Laosaurus*" *minimus*, Alberta, relationships (55).
Cretaceous, Lower: Foraminifera, new, Germany and England (5); molluscan faunas, fresh-water, North America, (87).
Cretaceous, Upper: elasmosaurid plesiosaur, South Dakota (80); Foraminifera, new, "*Clavulina*" shale, Peru (64).
Cricetid?, upper Eocene, Wyoming (83).
Crinoidea: *Jaekelicrinus bashkiricus* Yakovlev, n. gen., n. sp., upper Devonian, Russia (84).
Cruciloculina d'Orbigny, validity of (2).
Danian: superfluous term in Near East (74).
Dentalium (*Antalis*) *raymondi*, giant scaphopod, Pennsylvanian, Texas (46).
Devonian: *Microcyclus*, new species, Michigan and Ontario (65); nautiloids (21).
Devonian, upper: *Jaekelicrinus bashkiricus* Yakovlev, n. gen., n. sp., Russia (84).
Dictyoconus lineage: parallelism with rotaliid Foraminifera of the Chapmanininae (23).
Dinosauria: "*Laosaurus*" *minimus*, relationships, Cretaceous, Alberta (55).
Duchesnian: small mammals, Wind River Basin, Wyoming (83).
Dutch East Indies: coralline algae, Tertiary (38).
East Indies, Dutch: coralline algae, Tertiary (38).
Echinoidea: Cretaceous, Peru (12).
Ecuador: Cancellariidae, Miocene (45); *Hannatoma* fauna, age of (44, 61, 97).
Eden faunule, Martinsburg group, New Jersey, correlation with eastern Pennsylvania (81).
Egg capsules, chimaeroid fishes, Triassic, Connecticut Valley (6).
Egypt: Foraminifera, upper Tertiary (60); Globigerinidae ratios, Miocene (75).
Elasmosaurs: *Alzadasaurus pembertoni*, Upper Cretaceous, South Dakota (80).
Elvinia zone: trilobites, faunules, upper Cambrian, Texas (82).
England: new Foraminifera, Lower Cretaceous (5).
Eobronteidae: term to replace Scutellidae, Richter and Richter (57).
Eobronteus hibernicus: correction of location (56).
Eocene: coralline algae, Dutch East Indies (38); -Cretaceous boundary, established by quantitative generic analyses, Near East (74); "Eocene fresh-water Mollusca from Wyoming" Jour. Paleontology, Vol. 22, pp. 634–640, location corrections (85); Foraminifera and correlation of Cane River formation, Louisiana (36); *Hannatoma* fauna, Ecuador (61, 97), Peru (44, 97); Venezuela (39, 97); Ostracoda, Wyoming, Montana, Utah, Colorado (67).
Eocene, middle: *Ferayina*, Venezuela (10); gastropods, California (29).
Eocene, upper: Carbonera formation, Colombia, age of (18); Chira group, Peru, fauna (63); Foramin-

ifera, larger, Panama Canal Zone (11); Foraminifera, new, Talara formation, Peru (64); *Hannatoma* fauna, Colombia (97), Ecuador (61); Venezuela (97); small mammals, Duchesnian, Wyoming (83).
Eomyid, upper Eocene, Wyoming (83).
Errata: No. 4, p. 441; No. 6, p. 692.
Fellowships: National Research Council (95); Geology Department, School of Mines and Metallurgy, University of Missouri (96).
Ferayina, Foraminifera, Middle Eocene, Venezuela (10).
Foraminifera: Asagai formation (Tertiary), Japan (1); bibliography and index to new genera, species, and varieties, 1947 (70), 1948 (71), supplements and corrections, 1931–1947 (69), 1939–1947 (71); *Cibicides*, nomenclatural notes (62); Cretaceous, Lower, new genera, England and Germany (5); Cretaceous, Lower, Walnut formation, northern Texas and southern Oklahoma (41); Cretaceous, Upper, new genera, Peru (64); *Cruciloculina* (2); Eocene, Cane River formation, Louisiana (36), new names for two species, Clarke County, Alabama (4); Eocene, upper, larger, Panama Canal Zone (11); *Ferayina*, middle Eocene, Venezuela (10); fusulinids, Permian, Maracaibo Basin, Colombia and Venezuela (73), Permian, Timor (72), Permian, lower, new type, Japan (30); Globigerinidae ratios, value in stratigraphy, Miocene, Egypt and Turkey (75); *Hagenowella*, Paleocene, Netherlands (34); *Hannatoma* fauna, South America (19, 32, 39, 44, 61, 97); Kokozura formation, Miocene, Japan (3); *Lepidocyclina*, new, Miocene, Japan (50); Oligocene, Clarke County, Alabama, new names for two species (4); *Pullenia*, nomenclatural notes (62); Rotaliidae, Chapmanininae (10, 23, 24); Tertiary, upper, Egypt (60).
Franconian age, trilobites, Oklahoma (22).
Fusulinids: Permian, Maracaibo Basin, Colombia and Venezuela (73), Timor (72); Permian, lower, new type, Japan (30).
Gastropoda: Cancellariidae, nomenclatural units and tropical American species (45); Eocene, middle, new species, California (29); *Pedioceras*, synonym of *Crioceras* (*Pseudocrioceras*) (88).
Gatun Lake, Panama Canal Zone, larger Foraminifera from upper Eocene (11).
Genomorph concept: bearing of *Lithostrotion*, Mississippian, British Columbia, on (42).
Georgia: ostracodes, leperditiid (68).
Germany: new Foraminifera, Lower Cretaceous (5).
Globigerinidae: ratios, value in stratigraphy, Miocene, Egypt and Turkey (75).
Graptolites: new varieties, Ordovician, Tennessee and Virginia (17).
Gulf Coast: *Eucytherura*, Cretaceous and Tertiary (78).
Hagonowella, new species, Paleocene, Netherlands (34).
Hannatoma fauna: California (44); Colombia (19, 32, 97), Ecuador (44, 61, 97); Peru (44, 97); Venezuela (19, 32, 97).
Hannatoma zone: stratigraphic position (Eocene) in western Táchira region, Venezuela (39).
Herpetocypris Sars (67).
Hinge: features, leperditiid ostracods, Georgia (68); mechanism Ostracoda, Paleozoic (59).
Homonyms: in "Early Upper Cambrian faunas of Central Montana" (40); synonymous, possibility of (27).
Honey Creek formation (Upper Cambrian): redefinition of scope, trilobites (22).
Hoplophoneus, statistical study (35).
Howellaspis Lochman and Denson (40).
Illinois: cephalopods, Mississippian (47).
Index to Foraminifera: new genera, species, and varieties, 1947 (70), 1948 (71); supplements and corrections, 1931–1947 (69), 1939–1947 (71).
Insectivores: *Metacodon*, upper Eocene, Wyoming (83).
Iowa: cephalopod coquina, Ordovician (48); conodonts, Mississippian, late (92), Mississippian, lower (93), Pennsylvanian, early (91).
Ireland: Craighead limestone at Girvan, age of (56); *Eobronteus hibernicus*, County Tyrone (56).
Isopod track, *Oniscoidichnus*, Coconino sandstone (7).
Isopodichnus Brady (7).
Jaekelicrinus bashkiricus Yakovlev, s. sp., n. gen., upper Devonian, Russia (84).
Jamaica: Cancellariidae, Miocene (45).
Japan: *Cruciloculina*, Sawane formation (Pliocene) (1); Foraminifera, Asagai formation (Tertiary) (1); Kokozura formation (Miocene) (3), fusilinid, Lower Permian (30); new Mollusca, Tertiary (31).
Java: coralline algae, Tertiary (38).
Kentucky: cephalopods, Mississippian (47).
Lagena parvulipora (4).
Laosaurus minimus (55).
Leperditia hisingeri var. *gibbera*, *L. gigantea* (68).
Leperditiid ostracods: muscle marks, hinge and overlap features, new genera, modifications, Georgia (68).
Lepidocyclina: mortoni (11); Miocene, new, Japan (50).
Lithostrotion: Mississippian, British Columbia, bearing on genomorph concept (42).
Loganopeltis: facial sutures, synonym of *Loganopeltoides* (54).

Loganopeltoides: facial sutures, synonym of *Loganopeltis* (54).
Louisiana: Foraminifera from Cane River Eocene (36).
Maquoketa formation (Ordovician): cephalopod coquina, Iowa (48).
Maryvillia aequa (40).
Metacodon: upper Eocene, Wyoming (83).
Michigan: *Microcyclus*, new species, Devonian (65).
Microcyclus: new species, Devonian, Michigan and Ontario (65).
Miocene: Cancellariidae, tropical America (45); coralline algae, Dutch East Indies (38); Foraminifera, Japan (3); Egypt (60); Globigerinidae ratios, Egypt and Turkey (75); *Lepidocyclina*, new, Japan (50).
Mississippian: *Athyris*, Missouri (76); cephalopods, central and western United States (47); *Lithostrotion*, British Columbia, bearing on genomorph concept (42).
Mississippian, lower: conodonts, Iowa (93).
Mississippian, upper: cephalopods and correlation, White Pine shale, Nevada (89, 90); conodonts, Iowa (92).
Missouri: *Athyris*, Mississippian (76); cephalopods, Mississippian (47); Geology Department, School of Mines and Metallurgy, University of Missouri, fellowships and assistantships (96).
Mollusca: Cretaceous, Lower, fresh-water, North America (87); *Hannatoma* fauna, Tertiary, California (44), Colombia (19, 32, 97), Ecuador (44, 61, 97), Peru (44, 97), Venezuela (19, 32, 97); location corrections for "Eocene fresh-water Mollusca from Wyoming," Jour. Paleontology, Vol. 22, pp. 634–640 (85); scaphopod, giant, Pennsylvanian, Texas (46); Tertiary, Japan (31).
Montana: cephalopods, Mississippian (47); homonyms and synonyms in "Early upper Cambrian faunas of central Montana" (40); Ostracoda, Eocene, Paleocene (67).
Multivariate analysis (9).
Muscle marks: leperditiid ostracods, Georgia (68).
Mytilus, new species, Pliocene, California (43).
National Research Council, fellowshp (95).
Nautiloidea: Devonian, new genera (21); Permian, Colombia (73).
Nautilus beccarii Linne, nomenclature and generic position (24).
Near East: "Danian" (74).
Netherlands: *Hagenowella*, Paleocene Foraminifera (34).
Nevada: cephalopods, Mississippian (47); Mississippian, late, and correlation of White Pine shale (89, 90).
New Jersey: Eden faunule, Martinsburg group (81).
New names: Brachiopoda: *Raymondella* Whittington to *Bancroftina* Sinclair (58); *Reedaspis* Prantl and Přibyl to *Raymondella* Reed (58); Cephalopoda: *Pedioceras* Gerhardt to *Crioceras* (*Pseudocrioceras*) Karsten (88); *Scaphites morrowi* to *S. pygmaeus* (37); Dinosauria: *Laosaurus minimus* to "*L.*" *minimus* (55); Echinoidea: *Cottaldia australis* to *Orthopsis* (12); Foraminifera: *Lagena parvulipora* to replace *L. elliptica* (4); *Lepidocyclina mortoni* to *L.* (*L.*) *montgomeriensis* (11); *Triloculina pinguis* to *T. subrotunda* (4); Gastropoda: *Carinorbis* Yen to *Carinulorbis* Yen (86); Invertebrate tracks: *Isopodichnus* Brady to *Oniscoidichnus* Brady (7); Ostracoda: *Leperditia gigantea* to *Herrmannina welleri* (68); *Leperditia hisingeri* var. *gibbera* to *L. longigibbera* (68); *Herpetocypris* Sars to *Erpetocypris* Brady and Norman (67); Pelecypoda: *Chione cathcartensis* to *Macrocallista c.* (52); *C. columbiensis* to *Protothaca c.* (52); *C. fluctifraga* to *Protothaca f.* (52); *C. mathewsonii* to *Amiantis m.?* (52); *C. whitneyi* to *Dosinia w.* (52); *Venus* (*C.*) *conradiana* to *Clementia* (52); Trilobita: *Arapahoia convexa* to *A. polita* (40); *A. neihartensis* to *A. elongata* (40); *A. prolixa to A. polita* (40); *A. walcottae* to *A. snowiensis* (40); *Bolivina astoriensis* Rau to *B. chehalisensis* (No. 4, p. 441); *Coosia grandis* to *C. dakotensis* (40); *Howellaspis* Lochman and Duncan to *Bolaspidella* Resser (40); *Maryvillia aequa* to *M. wyomingensis* (40); *Pilgrima* Duncan to *Paracedaria* Duncan (40).
Niobrara formation (Upper Cretaceous): new rudistid, Colorado (26).
Nomenclature: comments on "Paleogeography and nomenclature" (51); *Pullenia* and *Cibicides* (62); revision of international rules for zoological (33).
Ohio: conodonts, Pennsylvanian (66).
Oklahoma: Foraminifera, Lower Cretaceous (41); trilobites, upper Cambrian (22).
Oligocene: Foraminifera, Egypt (60); *Hannatoma* fauna, California and Ecuador (44).
Ontario: *Microcyclus*, new species, Devonian (65).
Ordovician: cephalopod coquina, Iowa (48); graptolites, Tennessee and Virginia (17); trilobites (hypoparian), Quebec (53).
Ordovician, middle: *Eobronteus*, North America (57).
Ordovician, upper: Craighead limestone at Girvan, Ireland, age of (56); Eden faunule, Martinsburg group, New Jersey (81); *Eobronteus* from Europe (57).
Ornament: suggested replacement by term *prosopon* (25).
Ostracoda: *Eucytherura*, Cretaceous and Tertiary, Gulf Coast (78); leperditiid, muscle marks, hinge and overlap features, new genera, modifications of classification, Georgia (68); Paleozoic, hinge mechanism (59); Pliocene, southern California (13); Tertiary, non-marine, western interior United States (67).

Overlap features: leperditiid ostracodes, Georgia (68).
Pacific Coast of North America: *Chione* and *Securella*, Tertiary and Recent species (52).
Paleocene: *Hagenowella*, Netherlands (34); Ostracoda, Wyoming, Montana, Utah, Colorado, North Dakota (67).
"Paleogeography and nomenclature," comments on (51).
Paleontologist: biologist or geologist (53).
Paleontology: Center of Paleontological Studies and Documentation, Paris Museum (94).
Paleontology, quantitative: analysis of variance (49); applied to the determination of Cretaceous-Eocene boundary, Near East (74); Globigerinidae ratios, Miocene, Egypt and Turkey, value in stratigraphy (75); multivariate analysis (9); statistical study of *Hoplophoneus* (35).
Paleozoic: Ostracoda, hinge mechanism (59).
Panama Canal Zone: larger Foraminifera from Upper Eocene, Gatun Lake (11).
Pedioceras Gerhardt (88).
Pelecypoda: fossil and Recent species, Pacific Coast, North America (52); *Mytilus*, new species, Pliocene, California (43); new rudistid, Upper Cretaceous, Colorado (26).
Pella beds (Late Mississippian): conodonts, Iowa (92).
Pennsylvania: correlation of Martinsburg group, New Jersey, with formations in eastern (81).
Pennsylvanian: conodonts, Ohio (66).
Pennsylvanian, lower: conodonts, Iowa (91).
Pennsylvanian, middle: giant scaphopod, Texas (46).
Permian: fusulinids, Timor (72); fusulinids and cephalopods, Maracaibo Basin, South America (73); invertebrate track, *oniscoidichnus*, Coconino (7).
Permian, lower: fusulinids, Japan (30).
Peru: Cancellariidae, Miocene (45); Chira group, age of (63); Echinoids, Cretaceous (12); Foraminifera, new genera, Upper Cretaceous and Eocene (64); *Hannatoma* fauna, (44, 97).
Photography: stereophotography (20).
Pilgrima Duncan (40).
Plesiosaura: *Alzadasaurus pembertoni*, Upper Cretaceous, South Dakota (80).
Pliocene: *Cruciloculina*, Sawane formation, Japan (2); Foraminifera, Egypt (60); *Mytilus*, new species, California (43); Ostracoda, California (13).
Prosopon: proposed term to replace *ornament*.
Prospect Hill sandstone (Lower Mississippian): conodonts, Iowa (93).
Pullenia: nomenclatural notes (62).
Puma, fossil: *Hoplophoneus*, Tertiary, Wyoming (35).
Quantitative paleontology: analysis of variance (49); generic analysis of microfauna, use in establishing Cretaceous-Eocene boundary, Near East (74); Globigerinidae ratios, Miocene, Turkey and Egypt, value in stratigraphy (75); multivariate analysis (9); statistical study of *Hoplophoneus* (35).
Quebec: (hypoparian) trilobites, (Ordovician (53).
Rabbit, fossil: dental anatomy, upper Eocene, Wyoming (83).
Raymondella Whittington (58).
Recent: Pelecypoda, Pacific Coast (52).
Reedaspis, Prantl and Přibyl (58).
Reviews: Brachiopoda and Mollusca from the *Productus* limestones of the Salt Range, by F. R. Cowper Reed (8); Foraminifera in the Gullmar Fjord and the Skagerak, by Hans Höglund (14); Lower Ordovician Ostracods of the Siljan district, Sweden, by Ivar Hessland (77); Repartition stratigraphique du genre *Arenobulimina* Cushman, by N. Grekoff (15); The Swedish Paleocene and its foraminiferal fauna, by Fritz Brotzen (16).
Rodents, paramyd: Upper Eocene, Wyoming (83).
Rudistid, new: Upper Cretaceous, Colorado (26).
Russia: *Jaekelicrinus bashkiricus* Yakovlev, n. gen., n. sp., upper Devonian (84).
Scaphites morrowi, new name for *Scaphites pygmaeus* Morrow 1935 (37).
Scaphites pygmaeus (37).
Scaphopod, giant, Pennsylvanian, Texas (46).
Sciuravid: upper Eocene, Wyoming (83).
Securella: Tertiary and Recent, Pacific Coast of North America (52).
Silurian, basal: *Eobronteus*, Europe (57).
Society records and activities: Paleontological Society (no. 1, p. 105; no. 3, p. 331, p. 338; no. 5, p. 575); Society of Economic Paleontologists and Mineralogists (no. 2, p. 234; no. 4, p. 442, p. 449).
South America: *Hannatoma* fauna, age and environmental conditions (97).
South Dakota: elasmosaurid plesiosaur, Upper Cretaceous (80).
Species without a genus (28).
Spirifer whitneyi: analysis of variance applied to (49).
Sporobulimina: Foraminifera, new, Upper Cretaceous, Peru (63).
Sporobuliminella: Foraminifera, new Upper Cretaceous, Peru (63).
Statistical studies: analysis of variance (49), Globigerinidae ratios, Egypt and Turkey (75); *Hoplophoneus* (35); multivariate analysis (9).

Stereophotography, as paleontological tool (20).
Stichocassidulina: Foraminifera, new, upper Eocene, Peru (64).
Synonymous homonyms, possibility of (27).
Synonyms: in "Early upper Cambrian faunas of central Montana" (40).
Táchira region, Venezuela: *Hannatoma* fauna, stratigraphic position, Eocene (39).
Tetracoralla: new species, Devonian, Michigan and Ontario (65).
Talara formation (upper Eocene): new Foraminifera, Peru (64).
Tennessee: Ordovician graptolites (17).
Tertiary: coralline algae, Dutch East Indies (38); *Eucytherura*, Gulf Coast (78); Foraminifera, Asagai formation, Japan (1); *Hannatoma* fauna, Colombia and Venezuela (19, 32); *Hoplophoneus*, lower Brule, Wyoming (35); Mollusca, Japan (31); Pelecypoda, Pacific Coast (52).
Tertiary, early: ostracodes, non-marine, western interior United States (67).
Tertiary, upper: Foraminifera, Egypt (60).
Texas: Foraminifera, Walnut formation, Lower Cretaceous (41); scaphopod, giant, Pennsylvanian (46); trilobite faunules, Upper Cretaceous (82).
Timor: fusulinids, Permian (72).
Track, invertebrate: *Oniscoidichnus*, Permian (7).
Triassic: chimaeroid egg capsules, Connecticut Valley (6).
Trilobita: *Elvinia* zone, upper Cambrian, Texas, new genera, faunules (82); *Eobronteus hibernicus*, upper Ordovician?, Ireland (56); (hypoparian) trilobites, facial sutures and validity of *Loganopeltoides* and *Loganopeltis*, Ordovician, Quebec (53); *Raymondella Reed* 1935 (58); upper Cambrian, Oklahoma (22).
Triloculina pinguis (4).
Trinidad: Cancellariidae, Miocene (45).
Turkey: Globigerinidae ratios, Miocene (75).
Utah: Ostracoda, Eocene, Paleocene (67).
Variance, analysis of: application to paleontology (49).
Venezuela: *Ferayina*, middle Eocene (10); *Hannatoma* fauna, Tertiary (19, 32, 39); fusulinids, Permian (73).
Venus (Chione) conradiana (52).
Virginia: gaptolites, Ordovician (17).
Walnut formation (Lower Cretaceous): Foraminifera, Texas and Oklahoma (41).
White Pine shale (Late Mississippian): cephalopods and correlation, Nevada (89, 90).
Wichita Mountains: trilobites, upper Cambrian (22).
Wilberns limestone (Upper Cretaceous): trilobite faunules, Texas (82).
Wyoming: *Hoplophoneus*, lower Brule formation (35); mammals, small, uppermost Eocene, Wind River Basin (83); location corrections on "Eocene fresh-water Mollusca from Wyoming," Jour. Paleontology, Vol. 22, pp. 634–640 (86); ostracodes, Eocene, Paleocene (67).
Zapotal sands: *Hannatoma* fauna, Ecuador (61).
Zoological nomenclature: revision of international rules (33).